

# Elevated atmospheric CO<sub>2</sub> and humidity delay leaf fall in *Betula pendula*, but not in *Alnus glutinosa* or *Populus tremula* × *tremuloides*

Douglas Godbold · Arvo Tullus · Priit Kupper ·  
Jaak Sõber · Ivika Ostonen · Jasmin A. Godbold ·  
Martin Lukac · Iftekhar U. Ahmed · Andrew R. Smith

Received: 28 December 2013 / Accepted: 18 April 2014 / Published online: 21 May 2014  
© INRA and Springer-Verlag France 2014

## Abstract

• **Context** Anthropogenic activity has increased the level of atmospheric CO<sub>2</sub>, which is driving an increase of global temperatures and associated changes in precipitation patterns. At Northern latitudes, one of the likely consequences of global warming is increased precipitation and air humidity.

**Handling Editor:** Thomas Wohlgemuth

**Contribution of the co-authors** Douglas Godbold. Instigated the BangorFACE experiment, analysed data and co-wrote the paper  
Arvo Tullus. Instigated the FAHM experiment, collected and analysed data and co-wrote the paper  
Priit Kupper. Instigated the FAHM experiment and co-wrote the paper  
Jaak Sõber. Instigated and operated the FAHM experiment and analysed the weather data  
Ivika Ostonen. Collected and analysed fine-root data  
Andrew Smith. Collected and analysed data and wrote the original draft  
Martin Lukac. Analysed data.  
Iftekhar Ahmed. Collected leaves and carried out laboratory and data analysis  
Jasmin Godbold. Carried out statistical analysis and data interpretation

**Executive summary** The effects of both elevated atmospheric CO<sub>2</sub> and increased air humidity on autumn leaf fall were assessed using free air systems. Both factors delayed leaf litter fall in *Betula pendula*, but not in *Populus tremula* × *tremuloides* or *Alnus glutinosa*.

**Electronic supplementary material** The online version of this article (doi:10.1007/s13595-014-0382-4) contains supplementary material, which is available to authorized users.

D. Godbold (✉) · I. U. Ahmed  
Institute of Forest Ecology, Universität für Bodenkultur (BOKU),  
1190 Vienna, Austria  
e-mail: douglas.godbold@boku.ac.at

A. Tullus · P. Kupper · J. Sõber  
Department of Botany, Institute of Ecology and Earth Sciences,  
University of Tartu, Lai 40, 51005 Tartu, Estonia

I. Ostonen  
Department of Geography, Institute of Ecology and Earth Sciences,  
University of Tartu, Vanemuise 46, 51005 Tartu, Estonia

• **Aims** In this work, the effects of both elevated atmospheric CO<sub>2</sub> and increased air humidity on trees commonly growing in northern European forests were assessed.

• **Methods** The work was carried out under field conditions by using Free Air Carbon dioxide Enrichment (FACE) and Free Air Humidity Manipulation (FAHM) systems. Leaf litter fall was measured over 4 years (FACE) or 5 years (FAHM) to determine the effects of FACE and FAHM on leaf phenology.

• **Results** Increasing air humidity delayed leaf litter fall in *Betula pendula*, but not in *Populus tremula* × *tremuloides*. Similarly, under elevated atmospheric CO<sub>2</sub>, leaf litter fall was delayed in *B. pendula*, but not in *Alnus glutinosa*. Increased CO<sub>2</sub> appeared to interact with periods of low precipitation in summer and high ozone levels during these periods to effect leaf fall.

• **Conclusions** This work shows that increased CO<sub>2</sub> and humidity delay leaf fall, but this effect is species-specific.

**Keywords** Climate change · Free air CO<sub>2</sub> enrichment (FACE) · Free air humidity manipulation · Leaf fall · Ozone

## 1 Introduction

Anthropogenic activities since the industrial revolution have increased atmospheric CO<sub>2</sub> concentrations (IPCC 2013),

J. A. Godbold  
School of Ocean and Earth Science, National Oceanography Centre,  
Southampton, University of Southampton,  
Southampton SO14 3ZH, UK

M. Lukac  
School of Agriculture, Policy and Development, University of  
Reading, Reading, Berkshire RG6 6AR, UK

A. R. Smith  
School of Environment, Natural Resources and Geography,  
Bangor University, Gwynedd LL57 2UW, UK

leading not only to climate warming, but also to direct effect of elevated CO<sub>2</sub> on forest net primary productivity (NPP, Zak et al. 2011). In addition, climate change is predicted to increase precipitation at Northern latitudes (IPCC 2013), likely leading to an increase in air humidity. For example, in the Baltic region, climate change scenarios for the year 2100 predict an increase in air temperature (by 2.3–4.5 °C), precipitation (by 5–30 %) and cloudiness (by 2 %), but also higher wind speeds and vapour pressure (Kont et al. 2003). Studies investigating the impact of global environmental change on terrestrial ecosystems have identified a consistent pattern of phenological change in the Northern hemisphere (IPCC 2013). Analysis of normalised difference vegetation index (NDVI) remote sensing data gathered during 1985–1999 has revealed an 18-day extension of the growing season in Eurasia (Zhou et al. 2001). Multiple drivers have been shown to differentially influence plant phenophases, earlier bud break has been correlated with atmospheric warming and delayed senescence (Menzel et al. 2006) and interactions between temperature and elevated atmospheric CO<sub>2</sub> concentrations have been described (Taylor et al. 2008). The process of senescence is governed by developmental age, but also influenced by various integrated endogenous and environmental signals (Lim et al. 2007). Environmental factors influencing leaf senescence can be grouped into: (i) abiotic factors that include drought, nutrient limitation, extreme temperatures, and ozone-induced oxidative stress, and (ii) biotic factors including, pathogen infection or shading by other plants (Li et al. 2000). Endogenous factors influencing senescence include carbon source-sink relationships, phytohormones, particularly jasmonic (JA) and abscisic acid (ABA), ethylene and salicylic acid (SA). The aforementioned phytohormones initiate senescence through cellular signalling pathways in response to various abiotic and biotic stresses that promote the expression of senescence-inducing genes (Morris et al. 2000).

Elevated atmospheric CO<sub>2</sub> has been shown to increase long-term forest net primary productivity (Zak et al. 2011), if nutrients are not limiting (Leuzinger and Hätenschwiler 2013). However, studies of the effects of elevated atmospheric CO<sub>2</sub> on tree autumnal phenophase have produced conflicting results. For example, elevated CO<sub>2</sub> advanced senescence in two varieties of *Pinus ponderosa* (Houpis et al. 1988) and also in *Populus trichocarpa* (Sigurdsson 2001), yet delayed senescence of *Quercus myrtifolia* (Li et al. 2000) and *Populus* species grown in freely rooted field conditions during the AspenFACE and POPFACE studies (Taylor et al. 2008). At the DukeFACE experiment, however, no effect on leaf phenology was observed in *Liquidambar styraciflua* (Herrick and Thomas 2003).

Air water vapour content determines the vapour pressure difference between ambient air and leaf interior (VPD<sub>L</sub>), a gradient which drives the transpiration process of plant foliage. At higher relative humidity, both VPD<sub>L</sub> and transpirational

flux decrease, which has been demonstrated in the Free Air Humidity Manipulation (FAHM) experiment in both *Betula pendula* Roth and *Populus tremula* L. × *Populus tremuloides* Michx. in rainy summers when soil water content is not limiting in ambient conditions (Kupper et al. 2011; Tullus et al. 2012). It has been shown that elevated humidity diminishes nutrient supply to the leaves and photosynthetic capacity, altering foliar and fine-root properties and tree growth rate (Tullus et al. 2012; Hansen et al. 2013; Parts et al. 2013; Sellin et al. 2013). However, the effect of air humidity changes on leaf fall in trees has not been studied to date.

Natural autumnal senescence is regulated by the interaction of a number of factors including day length and temperature, nitrogen and water supply, as well as sink strength within the plant (Winger et al. 2006). Thus, changes in the timing of leaf senescence are governed by, amongst other factors, assimilation during the vegetation period and sugar accumulation in leaves (Swartzberg et al. 2010). Several studies utilising molecular genetic approaches have indicated that high concentrations of leaf sugars reduce photosynthetic activity, which in turn induces leaf senescence (Swartzberg et al. 2010). In *Acer saccharum*, girdling resulted in increased sugar accumulation in leaves and subsequent formation of anthocyanins (Murakami et al. 2008), whilst increased anthocyanin content in another study utilising the same species was associated with a delay in leaf senescence (Schaberg et al. 2008). Furthermore, transcriptome analysis of *Populus* trees grown under elevated CO<sub>2</sub> in field conditions revealed up-regulation of genes determining anthocyanin production during delayed senescence (Tallis et al. 2010). These authors suggest that anthocyanins may play a protective role in leaf metabolism and increase leaf longevity.

In the work presented here, we investigated the effect of two factors of global climate change, atmospheric CO<sub>2</sub> and humidity, on autumn leaf fall. We speculated the effects of both of these factors were tree species-specific. Thus, we hypothesised that (i) elevated CO<sub>2</sub> delays and (ii) elevated atmospheric humidity anticipates leaf senescence in broadleaved species.

## 2 Material and methods

The investigation was carried out at two sites, a Free Air Carbon dioxide Enrichment experiment (BangorFACE) and a Free Air Humidity Manipulation (FAHM) experiment.

### 2.1 The FACE facility

The BangorFACE experimental site was established in March 2004 on two former agricultural fields with a total area of 2.36 ha at the Bangor University research farm (53°14' N, 4°01' W) in North Wales, UK. Both fields were originally

pastures; one field was used for small-scale forestry experiments for the last 20 years, the other field was ploughed and planted with oil seed rape in 2003. Climate at the site is classified as Hyperoceanic, with a mean annual temperature in 2005 through 2008 of 11.5 °C and an annual rainfall of 1,034 mm (Fig. 1a). Soil is a fine loamy brown earth over gravel (Rheidol series) and classified as Fluventic Dystrochrept (Smith et al. 2013a). Soil texture is 63 % sand, 28 % silt and 9 % clay. The topography consists of a shallow slope of approximately 1–2° on a deltaic fan. The site aspect is northwesterly, with an altitude of 13 to 18 m a.s.l. The depth of the water table ranges between 1 and 6 m.

At the BangorFACE site, eight octagonal plots, four ambient and four CO<sub>2</sub> enriched were established, creating a 2×4 factorial block design across the two fields. Three tree species (*Alnus glutinosa* (L.) Gaertner, *Betula pendula* Roth. and *Fagus sylvatica* L.) were selected due to their contrasting shade tolerance, successional chronology and to represent a range of taxonomic, physiological and ecological types. Each plot was divided into seven planting compartments and planted in a pattern creating areas of one, two and three species mixtures. The present study makes use of observations originating from the single-species subplots of *B. pendula* and *A. glutinosa*. The site was planted with 60-cm saplings of each species. Within each treatment, the planting pattern was rotated by 90° between the four plots to avoid potential artefacts introduced by microclimate, soil and uneven growth rates of the different species. Each plot was surrounded by a 10-m border of *B. pendula*, *A. glutinosa* and *F. sylvatica* planted at the same density. The remaining field was planted at a 1-m hexagonal spacing with a mixture of birch (*B. pendula*), alder (*A. glutinosa*), beech (*F. sylvatica* L.), ash (*Fraxinus excelsior*

L.), sycamore (*Acer pseudoplatanus* L.), chestnut (*Castanea sativa* Mill.) and oak (*Quercus robur* L.). To protect the saplings, the entire plantation was fenced.


Carbon dioxide enrichment was carried out using high velocity pure CO<sub>2</sub> injection, with a target concentration in the FACE plots as ambient plus 200 ppm (Smith et al. 2013a). The elevated CO<sub>2</sub> concentrations, measured at 1-min intervals, were within 30 % deviation from the pre-set target concentration of 580 ppm CO<sub>2</sub> for 75–79 % of the time during the photosynthetically active part of 2005–2008 (Smith et al. 2013a). Vertical profiles of CO<sub>2</sub> concentration measure at 50-cm intervals through the canopy showed a maximum difference of 7 %.

Air temperature and precipitation were monitored using an automatic weather station (Campbell Scientific, Logan, UK) sampling at 3 m above the ground at hourly intervals. Ground-level ozone concentration was measured at a DEFRA air quality monitoring station at Aston Hill (52° 30' N, 3° 02' W) ca. 50 km from BangorFACE at hourly intervals and was matched to measurements made at the Centre for Ecology and Hydrology ozone research facility directly next to the BangorFACE site (53° 14' N, 4° 01' W).

## 2.2 FAHM facility

The Free Air Humidity Manipulation (FAHM) experimental facility is located at Järvselja Experimental Forest District in East Estonia (58° 14' N, 27° 18' E). The study area lies in the northern part of the temperate climate zone in the transition zone between maritime and continental climate. The study period comprised two growing seasons with drought conditions (2010 and 2011) and three with average precipitation

**Fig. 1** Monthly mean air temperature (line) and total precipitation (columns) at a BangorFACE during the years 2005–2008 and at b FAHM during the growing seasons 2008–2012



conditions (2008, 2009 and 2012) (Fig. 1b). Soil is classified as Endogleyic Planosol (Hansen et al. 2013). The FAHM site is a 2.7-ha fenced area, previously used for agriculture, where nine experimental circle plots are situated. Three experimental plots act as control plots. In three plots, the relative air humidity (RH) is elevated by approximately 7 % over ambient level using a misting technique (water is vaporised to a droplet size ca. 10  $\mu\text{m}$ ) and FACE-like technology to mix humidified air inside the plots (for more detailed technical description see Kupper et al. 2011 and Tullus et al. 2012). Humidification is applied when ambient RH < 75 %, air temperature > 10 °C and wind speed < 4 m/s. Three experimental plots were “open-top” plots from 2009 to 2011 and are not included in the current study. Half of each plot was planted with silver birch (*Betula pendula* Roth) and another half with hybrid aspen (*Populus tremula* L.  $\times$  *Populus tremuloides* Michx.) in 2006. The experimental plots are surrounded by a buffer zone, composed of hybrid aspen. Humidity manipulation experiment started in 2008 and has been running during all growing seasons (May–Oct) since then. The first experimental period with *B. pendula* ended in 2011; after that, a new birch generation was established with planted seedlings. Hybrid aspens were cut in 2012, and a new generation emerged as regrowth roots and stump sprouts.

Air temperature and precipitation were monitored using an automatic weather station (Campbell Scientific, Logan, UK) collecting in 10-min intervals at 6 m above the ground. Temperature data were collected in 10-min intervals. Winter precipitation (snow) data was obtained from the Estonian Environment Agency’s weather station, situated ca. 70 km from the FAHM site.

## 2.3 Litter collection

### 2.3.1 BangorFACE

Following observation of leaf fall, fallen leaf litter was collected at weekly intervals using litter baskets with an area of 0.11 m<sup>2</sup> until all leaves had abscised (September to December). A litter basket was located in each of the single species subplots. Litter was returned to the laboratory on the day of collection, washed and sorted into individual species, and then dried at 80 °C for 24 h. The dry weight of each species was determined and recorded for each species subplot within each ambient and elevated CO<sub>2</sub> plot. *F. sylvatica* was not used as senesced leaves remained attached until bud burst the following spring. Leaf retention was calculated by subtracting fallen litter at each sampling collection from the total fallen litter after all the leaves had abscised.

### 2.3.2 FAHM

Litter was collected from three controls (C) and three humidified (H) plots. Under both *B. pendula* and hybrid aspen, two

litter baskets (0.21 m<sup>2</sup>) per species were installed. Litter collection started in the end of July/beginning of August and continued in ca. 2-week interval until all leaves had abscised (usually by mid-November). Birch litter was collected during four experimental years (2008–2011); after that, the first generation of birch trees was harvested. *P. tremula*  $\times$  *tremuloides* litter was collected during 5 years (2008–2012), after which the first generation of aspen was removed. Litter samples were dried at 70 °C to constant weight, and dry mass of the samples was determined. Leaf retention was calculated as described above.

## 2.4 Data analysis

Generalised additive models (GAM) and generalised additive mixed models (GAMMs; Zuur et al. 2007; Wood, 2014) were used to describe the percentage change in remaining leaf mass at each collection date between ambient and treatment plots. Visual assessments of variograms and residuals vs. fitted values found weak evidence of temporal autocorrelation. However, as the time series consisted of <20 data points, it was more appropriate to model the variance structure, rather than the autocorrelation structure (Zuur et al. 2009). For *A. glutinosa* and *P. tremula*, data exploration indicated violation of homogeneity of variances as a result of differences between FACE rings and precipitation, respectively. As a result, we used a random effects model to model variability caused by the factor “ring” (for *A. glutinosa*) and the variable “precipitation” (for *P. tremula*). The additive (GAM; *B. pendula*) and additive mixed models (GAMM; *A. glutinosa*, *P. tremula*) were modelled with a binomial distribution and a logistic link function (Zuur et al. 2009). For both the FAHM and FACE analyses, the initial models of the GAMs and GAMMs included a smoother over “collection day” (s(days)), the factors “treatment” (ambient or elevated), “year”, as well as “precipitation” and “ozone” for the FACE analyses and “precipitation” and “temperature” for the FAHM analyses. To estimate the optimal amount of smoothing for each smoother, we used cross-validation (Zuur et al. 2009) and alternative models were compared using the Akaike information criterion (AIC). Once the optimal model was identified, the residuals were re-examined to ensure that model assumptions were met. Analyses were conducted in R (R Development Core Team 2014) and the “mgcv” library for additive (mixed) models (Wood, 2014).

## 3 Results

### 3.1 Environmental factors

At BangorFACE during the 4-year experiment period, in the summers of both 2006 and 2008, there were two consecutive

months with extremely low precipitation (Fig. 1a). These months were June and July in 2006 and May and June in 2008. In 2006, the highest summer temperature of the period under observation was reached. The highest temperature of 34.3 °C (Table 1) was recorded in July 2006 during a week-long period of very high temperatures. Accumulative ozone over the threshold of 40 ppb (AOT<sub>40</sub>) was highest during 2006, with daily peaks in excess of 210 ppb. In 2008, over the year, neither cumulative precipitation was very low nor was cumulative AOT<sub>40</sub> very high. However, during the low-rainfall months of May and June, 50 % of the total annual AOT<sub>40</sub> exceedance occurred and levels of over 170 ppb were reached. Based on the growing degree days (GDD) and maximum temperature, 2007 was the coolest of the 4 years (Table 1).

At the FAHM site, the 5-year experiment period also included two consecutive years with conditions of drought in the middle of the growing season, 2010 and 2011. The year 2010 was the warmest of the 5 years of the investigation, with ca. double the number of growing degree days compared to 2008 and 2012 (Table 2). The year 2011 was the driest year for plant growth as spring precipitation was low (Fig. 1b).

### 3.2 Leaf fall

At both the FAHM and the BangorFACE sites, based on weekly observations, the timing of bud burst was not affected by either elevated humidity or CO<sub>2</sub>, respectively. The autumn leaf fall at the FAHM site was modelled using a GAMM for *P. tremula*×*tremuloides* and a GAM for *B. pendula*. The curves of the measured data (Fig. 2) and the modelled data (Fig. 3) showed a high degree of agreement. In *P. tremula*×*tremuloides*, the *r*<sup>2</sup> for the GAMM fit was 97 %, and in *B. pendula*, the *r*<sup>2</sup> for the GAM fit was 95 %. At the FAHM site, different patterns of leaf fall were observed between *B. pendula* and *P. tremula*×*tremuloides* (Fig. 2). In *B. pendula*, fall began earlier and continued over an 8–9-week period, where as in *P. tremula*×*tremuloides*, ca. 80 % of the

leaves were lost within a 2-week period. In all study years, the leaf fall of *B. pendula* was significantly delayed (Fig. 3, Table 2, Online Resource 1) and slower in the increased humidity plots (*p*<0.0001), whilst such a consistent trend was not detected in *P. tremula*×*tremuloides* (*p*<0.0001). In 2010, in *P. tremula*×*tremuloides*, leaf fall was significantly earlier in the increased humidity plots (*p*<0.0001, Fig. 3). Generally, in hybrid aspen, leaf fall started later and lasted for a shorter period. In control plots, leaf fall of *B. pendula* began in the first half of August, whereas in the increased humidity plots, leaves started to fall almost 4 weeks later (Fig. 2). Litter fall dynamics in both *P. tremula*×*tremuloides* and *B. pendula* appeared to be dependent on annual weather conditions. Litter fall started earlier and more vigorously in the years 2010 and 2011 with dry summers (Fig. 1b). But *B. pendula* litter dynamics were also affected by increased humidity even in wet years (Figs. 1a, 2 and 3). However, in the modelled data, inclusion of the treatment factors temperature and precipitation did not improve the GAM, and both variables were removed during the backward selection procedure. The prolonged leaf retention in *B. pendula* meant that the time of 50 % leaf fall was reached ca. 21 days later in the increased humidity plots (Table 2). However, the duration to 100 % leaf fall did not differ between the ambient and humidity treatment.

At BangorFACE, a similar pattern of leaf loss was observed in *B. pendula* and *A. glutinosa*. Again, the curves of the measured data (Fig. 4) and the modelled data (Fig. 5) showed a high degree of agreement, with the exception of *B. pendula* in 2007. In *A. glutinosa*, the *r*<sup>2</sup> for the GAMM fit was 95 %, and in *B. pendula*, the *r*<sup>2</sup> for the GAM fit was 89 %. Inclusion of the factors temperature, precipitation and ozone did not improve the GAM or GAMM, and again, these variables were removed during the backward selection procedure. In *A. glutinosa*, in 2007, leaf loss was significantly earlier in both ambient and elevated atmospheric CO<sub>2</sub> compared to the other years (Figs. 4 and 5, Online Resource 1). In *A. glutinosa*,

**Table 1** Environmental variables and the lifespan of the leaf canopy (bud burst to final leaf fall) in *Betula pendula* at BangorFACE throughout the 4 years of CO<sub>2</sub> enrichment

Year	T <sub>min</sub> (°C)	T <sub>max</sub> (°C)	GDD (base 10 °C)	Rain (mm)	Ozone (AOT40)	Ambient CO <sub>2</sub> canopy lifespan (days)	Elevated CO <sub>2</sub> canopy lifespan (days)
2005	-3.5	27.0	1,910	726	9,058	201	201 (+0)
2006	-5.5	34.3	2,065	1,111	12,931	176	190 (+14)
2007	-3.3	24.3	1,672	705	3,783	172	172 (+0)
2008	-4.5	25.4	1,788	1,077	7,561	165	177 (+12)

The effect of elevated CO<sub>2</sub> on canopy lifespan is shown in parenthesis in days. T<sub>min</sub> and T<sub>max</sub> are based on the daily minimum and maximum temperatures

GDD growing degree days

$$\text{GDD} = \left( \frac{T_{\min} + T_{\max}}{2} \right) - 10$$

**Table 2** Environmental variables and the lifespan of the leaf canopy (bud burst to final leaf fall) at FAHM throughout the 5 years of relative humidity (RH) manipulation

Year	T <sub>min</sub> (°C)	T <sub>max</sub> (°C)	GDD (base 10 °C)	Rain (May–Oct) (mm)	<sup>a</sup> Total precipitation (mm)	Ambient RH canopy lifespan (days)		Elevated RH canopy lifespan (days)	
						50 % fallen	100 % fallen	50 % fallen	100 % fallen
2008	−17.1	30.8	619	502	853	Aspen: 156 Birch: 168	170 205	156 (+0) 177 (+9)	170 (+0) 205 (+0)
2009	−20.7	30.7	1,015	468	696	Aspen: 160 Birch: 145	190 211	160 (+0) 166 (+21)	190 (+0) 211 (+0)
2010	−27.6	36.9	1,321	387	828	Aspen: 151 Birch: 123	193 205	137 (−14) 163 (+40)	193 (+0) 205 (+0)
2011	−28.8	32	1,043	261	669	Aspen: 154 Birch: 141	178 192	154 (+0) 153 (+12)	178 (+0) 192 (+0)
2012	−31.3	32.9	753	339	756	Aspen: 140 Birch: <sup>−b</sup>	171 <sup>−b</sup>	140 (+0) <sup>−b</sup>	171 (+0) <sup>−b</sup>

The effect of FAHM on canopy lifespan is shown in parenthesis in days. T<sub>min</sub> and T<sub>max</sub> are based on the average annual minimum and maximum temperatures

GDD growing degree days

$$\text{GDD} = \left( \frac{T_{\min} + T_{\max}}{2} \right) - 10$$

<sup>a</sup> Total annual precipitation recorded by the Estonian Environment Agency's weather station, situated ca. 70 km from FAHM

<sup>b</sup> Birches were harvested in dormant season of 2011/2012

leaf fall was not significantly affected by elevated atmospheric CO<sub>2</sub> (Fig. 5, Online Resource 1). In contrast, in *B. pendula*, leaf fall was delayed by elevated atmospheric CO<sub>2</sub> in the years 2006 and 2008 based on the measured data (Fig. 4), and in all years based on the modelled data (Fig. 5, Online Resource 1). In 2006, litter collection was initiated on the 20th September (day 263). Under ambient CO<sub>2</sub>, 3 weeks later on the 11th October (day 283), 61 % of the *B. pendula* leaf canopy was still retained in the crowns. In comparison under elevated CO<sub>2</sub>, 80 % of the leaf canopy was still present in the crowns of the trees on the same date. Under elevated CO<sub>2</sub>, *B. pendula* still had 61 % of the total canopy 14 days later on the 25th October (day 298), thus extending the lifespan of the canopy (Table 1). In 2008, litter collection started on the 26th September (day 269), and by the 24th October (day 297) in the ambient plots, 96 % of the leaf canopy had fallen. Under elevated CO<sub>2</sub>, on the 24th October, 89 % of the canopy had fallen, and to reach a level of 96 %, a further 12 days were required.


#### 4 Discussion

Plant leaf senescence is a complex process predominantly influenced by environmental factors such as temperature, light, nitrogen availability and soil moisture. An example of this was seen in *A. glutinosa* where early leaf fall in 2007 occurred in the coolest of the 4 years. In addition, plant

physiological interactions which affect leaf senescence include phytohormones, leaf sugar content and source-sink status of the plant (Winger et al. 2006; Taylor et al. 2008). The data presented here show that elevated CO<sub>2</sub> and increased humidity both result in 2 to 3 weeks longer leaf retention in *B. pendula*. This effect was not seen in either *A. glutinosa* under elevated CO<sub>2</sub> or in hybrid aspen (*P. tremula* × *tremuloides*) under increased humidity. On the contrary, in 1 year, 2010, in *P. tremula* × *tremuloides* under increased humidity, leaf fall was earlier. However, the effect of elevated CO<sub>2</sub> on leaf retention in *B. pendula* also appears modified by interactions with other environmental factors, such as periods of drought, high temperature and high levels of ozone. Also in *P. tremula* × *tremuloides*, the shorter retention occurred in the warmest year (2010).

Plant growth in an elevated CO<sub>2</sub> atmosphere is often associated with increased accumulation of leaf starch and sugars, whilst leaf N is reduced (Ainsworth and Long 2005). Studies of *Arabidopsis* have demonstrated that leaf senescence can be induced by low N availability, and that N deficiency can result in leaf sugar accumulation (Pourtau et al. 2004). In support of this, leaf N of *Q. myrtifolia* in summer was lower under elevated CO<sub>2</sub> than under ambient CO<sub>2</sub>, but higher in autumn (Li et al. 2000). The higher autumn leaf N contents were related to delayed leaf fall. At BangorFACE, N contents of *B. pendula* and *A. glutinosa* leaves were not changed under elevated CO<sub>2</sub> (Smith et al. 2013a) during the summer, and in *B. pendula* in the autumn (Ferreira et al. 2010). No autumnal


**Fig. 2** Percentage leaf mass remaining in the canopy of birch (*Betula pendula*) and hybrid aspen (*Populus tremula* × *tremuloides*) grown at ambient humidity or increased humidity (FAHM). Data points show mean ± SE. *n* = 3



leaf N data are available for *A. glutinosa*. In contrast, N content in both *B. pendula* and hybrid aspen leaves were significantly lower in increased humidity plots in rainy summers (Tullus et al. 2012; Sellin et al. 2013). This indicates that in species under consideration, a change in leaf N status is not a common factor related to longer leaf retention. A generally consistent response to the process of leaf senescence is an increase in sugar content (Quirino et al. 2000). Complex interactions during sugar metabolism could help to explain these observations, which are supported by the results of a sugar maple (*Acer saccharum*) girdling experiment where leaf sugar

accumulation initiated the formation of anthocyanin, a molecule associated with delayed senescence (Murakami et al. 2008). Furthermore, using *Populus* spp., specific cDNA microarrays up-regulated gene expression of leucoanthocyanidin dioxygenase (LDOX) and dihydroflavonol reductase (DRF), two enzymes involved in the biosynthesis of anthocyanin were observed, in addition to increased autumnal leaf sugar accumulation (Tallis et al. 2010). At BangorFACE, *B. pendula* glucose and total soluble sugars leaf content were increased in leaves collected during 2006 under elevated CO<sub>2</sub>, whereas only the contents of glucose increased in *A. glutinosa* (Ahmed 2006).

**Fig. 3** Percentage leaf mass remaining in the canopy of birch (*Betula pendula*) and hybrid aspen (*Populus tremula* × *tremuloides*) grown at ambient humidity or increased humidity (FAHM). Model predictions (solid lines) and 95 % confidence intervals (dashed lines) are shown for leaf mass remaining over time for individual years in the ambient (black) and elevated (grey) humidity treatments




Cytokinins are known to delay leaf senescence (Yong et al. 2000), and usually, an excellent negative correlation between leaf cytokinin content and autumnal phenophase exists during senescence (Buchanan-Wollaston 1997). However, the physiology and biochemistry relating to the production of cytokinins and their interactions with senescence processes are

poorly understood. Many researchers consider cytokinins to be predominantly root-sourced plant hormones, which are translocated from the roots through the xylem (Dong et al. 2008). The supposition that cytokinin synthesis occurs primarily in roots was supported by the discovery of IPT-genes that control cytokinin synthesis in plants (Chang et al. 2003).



**Fig. 4** Percentage leaf mass remaining in the canopy of birch (*Betula pendula*) and alder (*Alnus glutinosa*) grown at ambient or elevated atmospheric CO<sub>2</sub> (FACE). Data points show mean  $\pm$ SE.  $n=4$




As elevated CO<sub>2</sub> has been shown to increase carbon allocation to roots and mycorrhizal symbionts (Iverson 2010), elevated CO<sub>2</sub> may also raise cytokinin production and subsequently increase leaf cytokinin concentrations. In the BangorFACE experiment, the leaf area index was not different between ambient and elevated CO<sub>2</sub> (Smith et al. 2013a), but the numbers of root tips in *B. pendula* were increased by 31 and 41 % in 2006 and 2008 under elevated CO<sub>2</sub>, and in *A. glutinosa*, a decrease or a 20 % increase were found in 2006 and 2008, respectively (Smith et al. 2013b). Similarly, under FAHM, in *B. pendula*, the root tip frequency per DW was 20 and 7 % higher in 2009 and 2010, respectively (Parts et al. 2013), and the number of root tips m<sup>-2</sup> was increased by 42 % compared to ambient in 2011 (Ostonen, unpublished), but no data are

available for hybrid aspen. A feedback mechanism involving a higher number of root tips and thus greater cytokinin production has the potential to explain the longer leaf retention under FACE and FAHM. An increase in fine root growth is a common feature in trees under elevated CO<sub>2</sub> and has been suggested to be due to high C allocation to roots, but also as a mechanism to increase nutrient uptake to meet the demand of increased aboveground growth (Smith et al. 2013a). Similarly, elevated humidity increased specific fine-root length (SRL) increase in *B. pendula* and was interpreted as a morphological adaptation leading to an increase in the absorptive area to facilitate nutrient uptake (Parts et al. 2013).

At BangorFACE, the years of longer leaf retention, 2006 and 2008, were characterised by periods of low precipitation

**Fig. 5** Percentage leaf mass remaining in the canopy of birch (*Betula pendula*) and alder (*Alnus glutinosa*) grown at ambient or elevated atmospheric CO<sub>2</sub> (FACE). Model predictions (solid lines) and 95 % confidence intervals (dashed lines) are shown for leaf mass remaining over time. In *Betula pendula* this is for the individual years in the ambient (black) and elevated (grey) CO<sub>2</sub> treatments. In *Alnus glutinosa* shown are the individual years with the treatments combined, as there are no treatment effects, but a significant difference between 2007 and the other years



for two successive months in the summer and high tropospheric O<sub>3</sub> concentration during this period. The physiological mechanisms behind this effect can only be speculated upon. Both O<sub>3</sub> (Yendrik et al. 2013) and elevated CO<sub>2</sub> (Eamus and Jarvis 1989) have been shown to reduce stomatal conductance, and thus reduce instantaneous leaf water loss. Further, as O<sub>3</sub> has been reported to directly contribute to earlier leaf senescence (Yendrik et al. 2013), lower stomatal conductance under elevated CO<sub>2</sub> may reduce O<sub>3</sub> exposure. Common to both FACE and FAHM is the potential to lower transpiration loss either through lower stomatal conductance (in FACE) or through lower water vapour pressure gradient (in FAHM). Higher water retention by the ecosystem throughout the growing season may lead to lower cumulative water stress in dry summers. Alternatively, the higher root biomass as discussed above may be beneficial in drier periods and also contribute to

lower cumulative water stress. However, it should also be noted that both *A. glutinosa* and *P. tremula* × *tremuloides* displayed varying leaf fall pattern compared to *B. pendula*.

## 5 Conclusions

Two separate experiments, one increasing atmospheric CO<sub>2</sub> whilst the other increasing air humidity, have both shown that deciduous tree species can respond to changing atmospheric conditions by prolonging their growing season. This effect, however, is not universal and appears species-specific. Further, the ability of trees to respond to changing atmospheric composition by retaining their foliage for longer may be modified by interaction with other factors. This research shows that the recently observed increasing duration of foliage

cover in forests may not only be an effect of increasing tropospheric temperature, but also be driven directly by changing atmospheric composition.

**Acknowledgments** The FAHM study was supported by the Ministry of Education and Science of Estonia (grant SF SF0180025s12) and by the EU through the European Social Fund (Mobilitas postdoctoral grant MJD 257) and the European Regional Development Fund (Centre of Excellence ENVIRON) and Project no. 3.2.0802.11-0043 (BioAtmos). The development of BangorFACE site infrastructure was funded by SRIF. We thank the Aberystwyth and Bangor Universities Partnership Centre for Integrated Research in the Rural Environment and the Forestry Commission Wales for financially supporting the running costs of the experiment. Andrew Smith was supported by the Sir Williams Roberts PhD Scholarship match funded by the Drapers' Company.

## References

- Ahmed IUMT (2006) Leaf decomposition of birch (*Betula pendula*), alder (*Alnus glutinosa*) and beech (*Fagus sylvatica*) grown under elevated atmospheric CO<sub>2</sub>. Dissertation, Bangor University
- Ainsworth EA, Long SP (2005) What have we learned from 15 years of free air CO<sub>2</sub> enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO<sub>2</sub>. *New Phytol* 165:351–371
- Buchanan-Wollaston V (1997) The molecular biology of leaf senescence. *J Exp Bot* 48:181–199
- Chang H, Jones ML, Banowitz GM, Clark DG (2003) Overproduction of cytokinins in petunia flowers transformed with P<sub>SAG12</sub>-IPT delays corolla senescence and decreases sensitivity to ethylene. *Plant Physiol* 132:1–10
- Dong H, Niu Y, Li W, Zhang D (2008) Effects of cotton rootstock on endogenous cytokinins and abscisic acid in xylem sap and leaves in relation to leaf senescence. *J Exp Bot* 59:1295–1304
- Eamus D, Jarvis PG (1989) The direct effects of increases in the global atmospheric CO<sub>2</sub> concentration on natural and commercial temperate trees and forests. *Adv Ecol Res* 19:1–55
- Ferreira V, Gonçalves AL, Godbold DL, Canhoto C (2010) Effect of increased atmospheric CO<sub>2</sub> on the performance of an aquatic detritivore through changes in water temperature and litter quality. *Glob Chang Biol* 16:3284–3296
- Hansen R, Mander Ü, Soosaar K, Maddison M, Löhmus K, Kupper P, Kanal A, Söber J (2013) Greenhouse gas fluxes in an open air humidity manipulation experiment. *Landsc Ecol* 28:637–649
- Herrick JD, Thomas RB (2003) Leaf senescence and late-season net photosynthesis of sun and shade leaves of overstory sweetgum (*Liquidambar styraciflua*) grown in elevated and ambient carbon dioxide concentrations. *Tree Physiol* 23:109–118
- Houpis JLL, Surano KA, Cowles S, Shinn JH (1988) Chlorophyll and carotenoid concentrations in two varieties of pine ponderosa seedlings subjected to long-term elevated carbon dioxide. *Tree Physiol* 4:187–193
- IPCC (2013) *Climate change 2013: the physical science basis*. Cambridge University Press, Cambridge
- Iverson CM (2010) Digging deeper: fine-root response to rising atmospheric CO<sub>2</sub> concentration in forested ecosystems. *New Phytol* 186:346–357
- Kont A, Jaagus J, Aunap R (2003) Climate change scenarios and the effect of sea-level rise for Estonia. *Glob Planet Change* 36:1–15
- Kupper P, Söber J, Sellin A, Löhmus K, Tullus A, Rääm O, Lubenetis K, Tulva I, Uri V, Zobel M, Kull O, Söber A (2011) An experimental facility for Free Air Humidity Manipulation (FAHM) can alter water flux through deciduous tree canopy. *Environ Exp Bot* 72:432–438
- Leutzinger S, Hätenschwiler S (2013) Beyond global change: lessons from 25 years of CO<sub>2</sub> research. *Oecologia* 171:639–651
- Li JH, Dijkstra P, Hymus GJ, Wheeler RM, Piastuch WC, Hinkle CR, Drake BR (2000) Leaf senescence of *Quercus myrtifolia* as affected by long-term CO<sub>2</sub> enrichment in its native environment. *Glob Chang Biol* 6:727–733
- Lim PO, Kim HJ, Name HG (2007) Leaf senescence. *Ann Rev Plant Biol* 58:115–136
- Menzel A, Sparks TH, Estrella N, Koch E, Aasa A, Ahas R, Alm-Kubler K, Bissolli P, Brasklavská O, Briede A, Chmielewski FM, Crepinsek Z, Cumel Y, Dahl A, Defila C, Donnelly A, Filella Y, Jateczak K, Mage F, Mestre A, Nordli O, Penuelas J, Pirinen P, Remisova V, Scheifinger H, Striz M, Susnik A, van Vliet AJH, Wielgolaski F-M, Zach S, Züst A (2006) European phenological response to climate change matches the warming pattern. *Glob Chang Biol* 12:1–8
- Morris K, A-H-Mackerness S, Page T, John F, Murphy AM, Carr JP, Buchanan-Wollaston V (2000) Salicylic acid has a role in regulating gene expression during leaf senescence. *Plant J* 23:677–685
- Murakami PF, Schaberg PG, Shane JB (2008) Stem girdling manipulates leaf sugar concentrations and anthocyanin expression in sugar maple trees during autumn. *Tree Physiol* 28:1467–1473
- Parts K, Tedersoo L, Löhmus K, Kupper P, Rosenvald K, Söber A, Ostonen I (2013) Increased air humidity and understory composition shape short root traits and the colonizing ectomycorrhizal fungal community in silver birch stand. *For Ecol Manag* 310:720–728
- R Core Team (2014) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>. Accessed 19 Mar 2014
- Quirino BF, Noh YS, Himelblau E, Amasino RM (2000) Molecular aspects of leaf senescence. *Plant Sci* 5:278–282
- Schaberg PG, Murakami PF, Turner MR, Heitz HK, Hawley GJ (2008) Association of red coloration with senescence of sugar maple leaves in autumn. *Trees* 22:573–578
- Sellin A, Tullus A, Niglas A, Öunapuu E, Karusion A, Löhmus K (2013) Humidity-driven changes in growth rate, photosynthetic capacity, hydraulic properties and other functional traits in silver birch (*Betula pendula*). *Ecol Res* 28:523–535
- Sigurdsson BD (2001) Elevated [CO<sub>2</sub>] and nutrient status modified leaf phenology and growth rhythm of young *Populus trichocarpa* trees in a 3-year field study. *Trees* 15:403–413
- Smith AR, Lukac M, Hood R, Healey JR, Miglietta F, Godbold D (2013a) Elevated CO<sub>2</sub> enrichment induces a differential biomass response in a mixed species temperate forest plantation. *New Phytol* 198:156–168
- Smith AR, Lukac M, Bambrick M, Miglietta F, Godbold DL (2013b) Tree species diversity interacts with elevated CO<sub>2</sub> to induce a greater root system response. *Glob Chang Biol* 19:217–228
- Swartzberg D, Hanael R, Granot D (2010) Relationship between hexokinase and cytokinin in the regulation of leaf senescence and seed germination. *Plant Biol* 13:439–444
- Tallis MJ, Lin Y, Rogers A, Zhang J, Street NR, Miglietta F, Karnosky DF, De Angelis P, Calfapietra C, Taylor G (2010) The transcriptome of *Populus* in elevated CO<sub>2</sub> reveals increased anthocyanin biosynthesis during delayed autumnal senescence. *New Phytol* 186:415–428
- Taylor G, Tallis MJ, Giardina CP, Percy KE, Miglietta F, Gupta PS, Gioli B, Calfapietra C, Gielen B, Kubiske MEM, Scarascia-mugnozza GE, Kets K, Long SP, Karnosky DF (2008) Future atmospheric CO<sub>2</sub> leads to delayed autumnal senescence. *Glob Chang Biol* 14:264–275
- Tullus A, Kupper P, Sellin A, Parts L, Söber J, Tullus T, Löhmus K, Söber A, Tullus H (2012) Climate change at Northern latitudes: rising

- atmospheric humidity decreases transpiration, N-uptake and growth rate of hybrid aspen. *PLoS One* 7:e42648
- Winger A, Purdy S, MacLean JA, Poutau N (2006) The role of sugars in integrating environmental signals during leaf senescence. *J Exp Bot* 57:391–399
- Wood S (2014) The mgcv Package v. 1.3-28, Mixed GAM Computation Vehicle with GCV/AIC/REML smoothness estimation. <http://cran.rproject.org/web/packages/mgcv/mgcv.pdf>. Accessed 19 Mar 2014
- Yendrik CR, Leisner CP, Ainsworth EA (2013) Chronic ozone exacerbates the reduction in photosynthesis and acceleration of senescence caused by limited N availability in *Nicotiana glauca*. *Glob Chang Biol* 19:3155–3166
- Yong JWH, Wong SC, Letham DS, Hocart CH, Farquhar GD (2000) Effects of elevated [CO<sub>2</sub>] and nitrogen nutrition on cytokinins in the xylem sap and leaves of cotton. *Plant Physiol* 124:767–779
- Zak DR, Pregitzer KS, Kubiske ME, Burton AJ (2011) Forest productivity under elevated CO<sub>2</sub> and O<sub>3</sub>: positive feedbacks to soil N cycling sustain decade-long net primary productivity enhancement by CO<sub>2</sub>. *Ecol Lett* 14:1220–1226
- Zhou L, Tucker CJ, Kaufmann RK, Slayback D, Shabanov NV, Myneni RB (2001) Variations in northern vegetation activity inferred from satellite data of vegetation index during 1981 to 1999. *J Geophys Res* 106:20069–20083
- Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM (2009) *Mixed effects models and extensions in ecology with R*. Springer Verlag, New York, LLC, p 574
- Zuur AF, Ieno EN, Smith, GM (2007) *Analysing Ecological Data*. Springer, New York, p 680